



ICMR BULLETIN

Vol. 46, No. 1-6

January-June 2016

In This Issue...

Biology & control of *Anopheles culicifacies* Giles 1901

Biology & control of *Anopheles culicifacies* Giles 1901

Malaria is a major public health problem in India. The epidemiology of malaria is complex on account of multiple vectors, vast and varied terrain, and a number of contextual determinants¹⁻³. In Southeast Asia Region of the World Health Organization, India alone contributes nearly 80 per cent malaria cases, and about 95 per cent of the population is estimated to be living at risk of malaria with large concentration of cases in forests and in the hilly and inaccessible terrains⁴. In 2013, the National Vector Borne Disease Control Programme (NVBDCP) reported 0.88 million microscopically confirmed malaria cases and 440 deaths, and each year these figures are on the decline, although scientific institutions have repeatedly highlighted gross under-reporting of malaria morbidity and mortality⁴⁻⁷. *Plasmodium vivax* and *P. falciparum* are the two main malaria parasites that produce nearly equal proportion of cases (50:50), with a few cases of *P. malariae* from certain parts of Odisha (formerly Orissa), and sporadic reports of *P. ovale*. There are six major mosquito vectors in India (*Anopheles culicifacies*, *An. fluviatilis*, *An. stephensi*, *An. minimus*, *An. dirus*, and *An. sundaicus*), and all taxa except *An. stephensi* are species complex⁸. Among these, *An. culicifacies sensu lato* is the most abundant vector. It is widely distributed in rural and peri-urban India. *An. culicifacies* has played a major role in perennial malaria transmission in forested areas (e.g. Madhya Pradesh), and this vector species has penetrated deforested areas of Northeast⁹⁻¹². *An. culicifacies* is responsible for epidemic malaria in its range of distribution and may cause intense malaria transmission in an estimated 75 million tribal population. *An. culicifacies* is multiple resistant to insecticides, and its control alone costs about 80 per cent of budget year marked for malaria control of the NVBDCP of the Government of India⁴. Furthermore, *An. culicifacies* comprises five sibling species with varying responses to insecticides and transmission potential¹³. Resurgence of malaria in late 1960s in India and adjoining countries of South Asia was largely

Editorial Board

- Chairman** : **Dr. Soumya Swaminathan**
Director-General, ICMR
and Secretary, DHR
Ministry of Health and
Family Welfare,
Government of India
- Editor** : **Dr. V.K. Srivastava**
- Members** : **Dr. Rashmi Arora**
: **Dr. D.K. Shukla**
: **Dr. Vijay Kumar**
: **Dr. Chandrashekar**
: **Dr. R.S. Sharma**
- Publisher** : **Shri J.N. Mathur**

the result of failure in the control of *An. culicifacies*^{14,15}. The present review provides comprehensive information on *An. culicifacies*, the most important malaria vector in India and the neighbouring countries.

Taxonomy & distribution

An. culicifacies Giles 1901 belongs to subgenus *Cellia* and series *Myzomyia*. Three synonyms of *An. culicifacies* are reported from India. These are the *indica* Theobald, 1901 of Chennai (formerly Madras); *listonii* Giles, 1901 from Ellichpur, Maharashtra, and *punjabensis* James, 1911 from Punjab¹⁶. It is in the last three decades that cytogenetic research has revealed the presence of five sibling species of *An. culicifacies* provisionally designated as A, B, C, D and E¹⁷. These sibling species differ in their biology, vectorial capacity and response to malaria control interventions. *An. culicifacies* s.l. is widely distributed in arid and semi-arid zones ranging from Afghanistan, Bangladesh, Cambodia, Southern China, Eritrea, India, Iran, Laos, Myanmar, Nepal, Pakistan, Sri Lanka, Thailand, Vietnam and Yemen (Fig. 1).

Seasonal prevalence

An. culicifacies s.l. is among the most widely distributed mosquito species in India and occurs in all mainland zones including Kashmir and high elevation in the Himalayas excluding islands of Andaman &

Nicobar and Lakshadweep (Fig. 2). It is predominantly a vector of unstable malaria in the entire semi-arid and arid zones. Populations of *An. culicifacies* are low to scanty at higher altitudes and may transmit malaria under favourable conditions, e.g. global warming may bring some ecotones under transmission. It has been reported from 1000 to 2000 meters above mean sea level in Nilgiris hills and Kashmir, respectively¹⁸⁻²¹. This mosquito thrives well in plains receiving fair amount of rainfall; heavy rains in various parts of the country (e.g. Maharashtra, Uttarakhand), droughts and floods (e.g. Uttar Pradesh, Bihar), and excessive cold and hot weather in the plains are changing the ecology leading to re-distribution of vector densities capable of malaria transmission. Generally, *An. culicifacies* populations peak in monsoon and post-monsoon months, and from negligible numbers attain enormously high densities within 4-6 wk resulting in focal to regional epidemics^{9-11, 22}. It is an important vector of epidemic malaria throughout its range of distribution, although it is a rare species in the Western Ghats.

Larval ecology

An. culicifacies s.l. is a prolific breeder and its preferred habitats are numerous and varied^{16,18}. Irrigation channels, seepage, unused wells, field channels, waste irrigation water, hoof marks and cart tracks, etc. are some the preferred breeding places. *An. culicifacies*




Fig. 1. Distribution map of *An. (Cellia) culicifacies* species complex in the world. Red and blue color depicts the high and low probability of occurrence of this species. Black dots display the sites of data collected (Source: http://www.map.ox.ac.uk/client_media/medialibrary/2011/10/vector_Asia-Pacific_paper.pdf).




Fig. 2. Predictive map of the occurrence of *An. (Cellia) culicifacies* in India. Red and blue color depicts the high and low probability of its occurrence. (Source: <http://www.maps.ox.ac.uk/browse-resources/vector-occurrence/culicifacies/IND/>).

larvae have also been recorded in varying densities in pools, fallow fields, mining pits, river-bed pools, seaside marsh, turf pools, river edges, perennial streams, tanks, and artificial containers. The most suitable places for the breeding of *An. culicifacies* are small rocky pools and pits of perennial stream with clear water or with perceptible flow without shade and growth of any vegetation or macroscopic algae; and the least suitable

places have highly turbid, stagnant and/or brackish water with good growth of vegetation including floating, sub-merged or vertical vegetation, growth of blue green algae and rich plankton; and the intermediate level of preference is found in fields, and channels with flowing water. *An. culicifacies* preferentially breeds in sunlit water collections. It breeds profusely in the freshly dug out pits, ornamental waters, unused swimming pools, and rain filled borrow pits along side of railway tracks. Although *An. culicifacies* is believed to breed in fresh water bodies, but it has adapted to lay eggs and undergo pre-imaginal development in saline/brackish water²³.

Sibling species complex

An. culicifacies s.l. was first colonized in the laboratory by the National Institute of Malaria Research (formerly Malaria Research Centre) in 1977²⁴. Initially considered to comprise biological races it has now been characterized as species complex with five informally designated species A, B, C, D and E^{13,17}. Techniques are now available to identify these sibling species within the taxon *Anopheles culicifacies s.l.* (Table I). Among these, the use of diagnostic fixed readable inversions in the ovarian nurse cell's polytene chromosomes unequivocally helps to identify different sibling species. Initially, two sibling species A and B were identified near Delhi; one with standard arrangement in polytene X-chromosome (X^{+a+b}) designated as 'A' and the other with two paracentric inversions 'a' and 'b' with chromosome arrangement (Xab) designated as 'B', without any evidence of heterozygotes in natural populations²⁵. Subsequently, examination of polytene chromosomes of populations from other regions of India revealed presence of two fixed inversions, *i.e.* g^1 and h^1 in chromosome arm 2 in species B, and two distinct populations with chromosome arrangement

Table I. Techniques applied in the identification of *An. culicifacies* sibling species

Sibling species	Polytene chromosome inversion genotype	Mitotic karyotype Y-chromosome	LDH enzyme alleles	Cuticular hydrocarbon profile	Species specific DNA probes	PCR-RFLP	ASPCR
A	$X^{+a+b}; 2+g^1+h^1; +i^1/i^1$	Submetacentric	Fast	Identified	Yes	Yes	Yes
B	Xab; $2g^1+h^1$	Acrocentric, Submetacentric	Slow	Identified	Yes	Yes	Yes
C	Xab; $2+g^1+h^1$	Acrocentric, Submetacentric	Slow	Identified	As B	As B	Yes
D	$X^{+a+b}; 2i^1+h^1$	Submetacentric	Fast	Not done	Not tested	As A	Yes
E	Xab; $2g^1+h^1$	Submetacentric	Slow	Not done	Not tested	Yes	Yes

Source: Ref. 17 (reproduced with permission). PCR-RFLP, polymerase chain reaction-restriction fragment length polymorphism; ASPCR, allele specific polymerase chain reaction; LDH, lactate dehydrogenase

Xab; $2g^1+h^1$ and that with Xab; $2+g^1h^1$ were identified which were considered reproductively isolated. The new population with Xab; $2+g^1h^1$ chromosome arrangement was designated as species 'C'²⁶. Additionally, examination of a few more populations from north India revealed polymorphism for yet another inversion 'i' in chromosome 2 of species A which was observed to be fixed in populations from south India. The evidence of deficiency of heterozygotes in northern and central Indian populations for inversion 'i' and total absence of heterozygotes in populations of species A with X^{+a+b} ; $2+g^1+h^1$ and with X^{+a+b} ; $2i^1+h^1$ chromosome arrangement in southern populations provided cytogenetic evidence for species 'D' (X^{+a+b} ; $2i^1+h^1$)²⁷⁻²⁹.

In addition to sibling species A, B, C and D characterized by analyses of polytene chromosome arrangements, sibling species E was identified by Y-chromosome polymorphism in sibling species B populations from Rameshwaram island of Tamil Nadu, south India³⁰. Population with acrocentric Y-chromosome was retained as B while that of submetacentric Y-chromosome karyotype and high sporozoite infectivity was designated as new species E. Both sibling species B and E are homosequential for polytene chromosomes and occur in sympatry but apparently reproductively isolated by pre-mating isolating mechanisms.

Populations of *An. culicifacies* in range of its distribution other than India have also been examined for sibling species characterization, e.g. in Yeman and Iran, species A was identified^{31,32}, and both A and B are reported occurring sympatrically in Pakistan, of which species A was incriminated³³. In Sri Lanka, based on polytene chromosome analysis and DNA probes only species B was identified^{34,35}, which later confirmed to comprise another sibling species E, the one that was responsible for malaria transmission with high sporozoite infectivity³⁶. In Thailand, both A and B were identified³⁷.

The pre-mating reproductive isolation between sibling species was further substantiated by post-mating isolation mechanisms marked by unidirectional fertility/hybrid male sterility/ reduced fertility/ atrophied reproductive organs in reciprocal crosses between species A/B, and A/C. Reciprocal crosses between species B and C, however, produced fertile F1 hybrid males and females^{33,38,39}. In addition to these methods, these siblings could also be identified with reasonable accuracy by mitotic karyotype Y-chromosome



Fig. 3. Map of India showing geographical distribution of *An. culicifacies* complex (sibling species A, B, C, D, and E), and stratification (Divisions I-VII) for vector control options. (Source : Ref. 57).

polymorphism⁴⁰⁻⁴³, electrophoretic variation in lactate dehydrogenase enzyme for fast and slow allele (LdhF & LdhS)⁴⁴, cuticular hydrocarbon profiles⁴⁵, and highly repetitive DNA sequences⁴⁶. PCR-based diagnostic assays have been developed for sequencing 28S-D3 domain⁴⁷, ITS2-PCR-RFLP (restriction fragment length polymorphism)⁴⁸, rDNA-ITS2-PCR⁴⁹, which grouped *An. culicifacies* sibling species into two distinct groups namely group I (species A/D) and group II (species B/C/E). In another two step PCR assay based on sequence difference within the COII region, A/D specific primers distinguished species A and D, and B/C/E specific primers distinguished B, C and E⁵⁰, however, this assay could not be used to distinguish species B and E in Sri Lanka⁵¹. An efficient and less expensive multiplex PCR-based diagnostic assay using D2 domain of 28S rDNA has been reported which can consistently and accurately discriminate members of the species complex forming two unambiguous monophyly clades of species A/D (group I) and species B/C and E (group 2) were supported by strong bootstrap values⁵².

Sibling species distribution and sympatricity

Populations of *An. culicifacies* of diverse origin have been studied for distribution and abundance of its sibling species, A, B, C, D and E in India¹⁷. Among these, species B is the most predominant spread throughout the country and occurs sympatrically with

A or C or D (Fig. 3). Species A and B are sympatric in north and south India with predominance of species A in the north and species B in the south. In eastern Uttar Pradesh, north Bihar and northeastern States, species B is either predominant or the only prevalent species. Species B and C are predominant in the western and eastern regions. Species D is sympatric with species A and B in northwestern region, and with species A, B and C in central India and a few areas in southern India. Species E is sympatric with species B in southern Tamil Nadu including Rameshwaram islands and Sri Lanka, and there are reports of expansion of distribution range of species A, D and E in Odisha and that of species E in Madhya Pradesh but these investigations lack the component of chromosome analyses for diagnostic confirmation⁵³⁻⁵⁵. The proportions of sibling species, however, varied in different geographical zones and seasons, e.g. in Delhi where species A and B are sympatric, A was predominant throughout the year but proportions of B increased in post-monsoon months⁵⁶. In Alwar (Rajasthan), amongst sibling species A, B, C and D, species B increased in post-monsoon months, proportions of D remained the same throughout the year and densities of species C remained very low⁵⁷.

Bionomical characteristics

Five sibling species spread across India have distinct biological characteristics and role in malaria transmission (Table II). All members of *An. culicifacies* complex except E were predominantly zoophilic¹³. Species A had relatively high anthropophilic index (0-4%) compared to species B and D (0-1%), species C had intermediate level of anthropophilic index (0-3%), and species E, had the highest anthropophilic index (80%). All member species largely rest indoors human dwellings preferentially on roof ceilings after feeding on cattle, but also rests outdoors¹⁸. All are night biting species with different peak biting activity. The biting activity of A, B and C was observed all through the night except for D for which there was no biting after midnight. The peak biting activity of species A and B occurred between 2200 till 2300 h whereas for species C, it was seasonal; in April it occurred between 1800 till 2100 h and shifted to second quarter of the night in December¹⁷.

An. culicifacies s.l. has been incriminated by detection of gut and salivary gland infections by numerous independent investigators across its range of distribution throughout India^{16,18}. However, the seasonal infection rate varied from moderate to high in northwestern States, low to moderate in Deccan plateau,

low rates in Gangetic plains and east central region, and very low rates in northeastern States¹⁸. The sporozoite infectivity rates also varied among sibling species. Immunoradiometric based investigations revealed that sibling species A, C and D are vectors of *P. vivax* and *P. falciparum* (including drug resistant strains), and cumulative sporozoite infection rates were recorded 0.51, 0.3 and 0.4 per cent, respectively^{58,59}. Species B is a non-vector or poor vector evidenced by the low prevalence of malaria where species B is predominantly prevalent such as in eastern districts of Uttar Pradesh and southern Indian States⁶⁰. Species E is the most efficient vector and maintains endemic to epidemic malaria. These observations were further supported by comparative reproductive fitness for which sibling species B was observed to be less fit than species A and C of the complex as well as susceptibility to malaria parasite development⁶¹. Species A is susceptible to sporogony marked by higher oocyst and sporozoite rate than C and B; in species B parasite development is inhibited by oocyst encapsulation mechanism^{62,63}. Variations to development of insecticide resistance have also been reported among sibling species but with slower development of resistance to DDT in species A than species B and C, and faster development of resistance to malathion in C than B and at slower rate in A than B (Table II)^{64,65}.

Vector control

Indoor residual spraying (IRS) is the main stay for vector control in the country. For control of *An. culicifacies*, currently three rounds of spraying of malathion (25%WP, 2 g/m²) are undertaken in DDT resistant areas. In areas with double resistance, i.e. DDT and malathion, two rounds of synthetic pyrethroid insecticides, i.e. deltamethrin (2.5%, 20 mg/m²), or cyfluthrin (10% WP) or lambda-cyhalothrin (10% WP) or alphacypermethrin (5% WP) or bifenthrin (10% WP) each are sprayed as 25 mg/m². Malathion and/or synthetic pyrethroids are also sprayed to control epidemic malaria in complex emergency situations. In urban and industrial areas 5 per cent malathion thermal fogging is undertaken to contain build up of vector population.

The understanding of genetics of *An. culicifacies* species complex for its sibling species composition, distribution and behavioural characteristics has helped develop stratification for optimizing control efforts and saving costs (Fig. 3). Taking cognizance of the biological differences of sibling species, the rural India is stratified into seven divisions (excluding urban

Table II. Bionomical characteristics of *An. culicifacies* sibling species and role in malaria transmission in India

Bionomical characteristic	Sibling species				
	A	B	C	D	E
Anthropophilic index (%)	0-4	0-1	0-3	0-1	80
Biting activity (Peak biting activity in h)	All night (2200-2300)	All night (2200-2300)	All night (1800-2100)	Till midnight (1800-2100)	No data
Vector potential	Moderate	Poor	Moderate	Moderate	High
Sporozoite infection rate (%)	0.51	0.04	0.3	0.4	4.6
Breeding preferences	Rainwater, clean irrigation water	Riverine ecology	Rainwater, clean irrigation water	Rainwater, clean irrigation water	Riverine ecology
Insecticide resistance					
DDT	Slow (9-10 yr)	Fast (4-5 yr)	Fast (4-5 yr)	No data	No data
Malathion	Slow (9-10 yr)	Medium (6-7 yr)	Fast (4-5 yr)	No data	No data
Pyrethroid	No data	Medium (6-7 yr)	Medium (6-7 yr)	No data	No data

Source: Ref. 8 (reproduced with permission)

metropolitan cities) for benefit of prioritizing vector control options (Table III)⁵³⁻⁵⁷.

No specific strategy has been proposed for species D for its limited distribution and occurrence in low numbers often in sympatricity with other sibling species. In areas with predominance of species E like Rameshwaram Island and Odisha existing interventions should continue. In areas where A, C, and D are sympatric, the choice of insecticide should be guided by the susceptibility status of the predominant sibling species. In addition to stratification, it is strongly advised to develop stratification at the district/ block level to economize field operations. For example, in Shankargarh block of Allahabad district, Uttar Pradesh, the prevalence of malaria had direct correlation with cumulative abundance of sibling species A (64%) plus C (11%)⁶⁶. In general, it is proposed to integrate bioenvironmental methods; larvivorous fish (Guppy and Gambusia) in particular which have been applied successfully in different ecological terrains for control of vector breeding in malaria endemic States Gujarat, Maharashtra and Karnataka⁶⁷⁻⁷⁰.

Challenges in malaria control

India has reported appreciable decline in malaria cases from two million in 2001 to less than one million in 2013 and notable decline in malaria deaths⁴. However, there is a steady rise in *P. falciparum* proportions with large concentration of cases in tribal populations/forest

belts, hilly and difficult/ inaccessible areas. Malaria transmission is perennial affecting all age groups and with serious consequences to high risk groups, particularly with heavy death toll in pregnant mothers and infants^{4,7,71}. *An. culicifacies* transmitted malaria is highly uneven, maintains unstable to intermediate stability and brings periodical epidemics. This vector is responsible for malaria resurgence in India, Pakistan and Sri Lanka, although malaria is at the verge of elimination in Sri Lanka^{14-15,72-74}.

An. culicifacies maintains endemic malaria throughout rural India and its control has become a formidable task consuming huge resources year after year. Malaria control is particularly problematic in forested and degraded forests as in some areas spraying is difficult and remains unsupervised. Malaria transmission in forests is usually prolonged; for example, *An. culicifacies* was incriminated for over 10 months in a year in Balaghat area in Madhya Pradesh^{75,76}. *An. culicifacies s.l.* has grown multi-resistant to DDT, HCH (hexachlorocyclohexane) and malathion, in most parts of the country and in some States increased resistance to pyrethroids⁷⁷⁻⁸⁰. Molecular characterization revealed a low frequency of the *kdr* allele (mostly in heterozygous condition) in field populations that were resistant to DDT and pyrethroids^{81,82}. This species is invasive and expanding its territory by penetrating degraded forests of northeastern States of the country, formerly domain of *An. minimus* and *An. dirus*^{11,12}. In addition, global

Table III. Distribution of *An. culicifacies* sibling species and stratification for vector control options in India

Division	Distribution of sibling species and vector control options
I	No routine vector control measures are required except for chemotherapy of imported malaria. It includes States of Jammu and Kashmir, Himachal Pradesh, Punjab, Haryana, Delhi and parts of north-western Uttar Pradesh and Rajasthan. Sibling species A and B are sympatric; species B is predominant in northern region whereas in canal irrigated areas species A is in the majority.
II	Insecticide spray should be based on susceptibility status of species A or C. It includes States of Gujarat and districts of north-western Rajasthan, western Madhya Pradesh and north-western Maharashtra. Species B and C are widely prevalent except in irrigated areas where A is predominant. <i>An. culicifacies</i> is reported resistant to DDT and malathion.
III	No routine vector control measures are required except for chemotherapy of imported malaria. It includes districts of eastern and southern Uttar Pradesh, and districts of northern Bihar. In this division, species B is the only prevalent species. This division is less receptive to malaria transmission and measures such as minor engineering works for source reduction, use of biological control agents can be exploited for sustainable control.
IV	Indoor residual spraying (IRS) and existing interventions including insecticide-treated nets/ long-lasting insecticidal nets should continue. It includes seven States of northeastern region, districts of northern Bengal and State of Sikkim. In this division, <i>An. minimus</i> and <i>An. baimaii</i> are the two predominant vectors species; both are highly susceptible to DDT. <i>P. falciparum</i> is the predominant infection. Species B is the only member species of <i>An. culicifacies</i> but has very little role in malaria transmission.
V	Insecticide spray based on susceptibility status of species A or C should be applied. It includes most districts of Madhya Pradesh, north-western Maharashtra, northern Andhra Pradesh and districts of southern Uttar Pradesh. Transmission is intense and <i>P. falciparum</i> is the predominant infection. <i>An. culicifacies</i> is resistant to DDT and malathion, thus susceptibility status should be monitored for choice of insecticide targeting the most predominant sibling species. There are reports of invasion of species E in this zone warranting constant vigil and intensive control interventions to avert epidemics ⁵⁵ .
VI	Insecticide spray based on susceptibility status of species C and use of insecticide-treated nets/ long-lasting insecticidal nets are recommended. It includes State of Odisha (formerly Orissa), Chhattisgarh, Jharkhand, districts of southern Bihar, north-eastern districts of Andhra Pradesh. However, with recent reports of invasion of sibling species A, D and E, there is need for monitoring relative abundance for planning intensification of control interventions against A and E ^{53,54} .
VII	Insecticide spray and wherever feasible larvivorous fish should be applied. It includes districts of southern Andhra Pradesh, States of Karnataka, Kerala and Tamil Nadu. This division needs to be further stratified for distribution range of species A, B, C and E; the latter presently recorded from Rameshwaram island of Tamil Nadu. The control measures should be judiciously intensified against species A and E. The application of larvivorous fish has been a successful strategy in the control of breeding in wells.

Source: Refs 53-57.

warming, climate change, vector ecology, population migration, deforestation, developmental projects, and poor infrastructure have led to the opportunities for vector proliferation and increased malaria receptivity in the country^{83,84}.

The main strategy for malaria control in rural India continues to be indoor residual spraying (IRS) of insecticide based on the vector susceptibility status. It is the need of time to develop approaches for management of insecticide resistance for increased duration of its efficacy against target vector species by insecticide rotation, mosaic application, and integrating bio-environmental approaches⁸⁵⁻⁸⁸. IRS has become less effective and operationally difficult on account of poor acceptance by communities⁸⁹. Instead of IRS,

implementation of insecticide-treated netting materials / long-lasting insecticidal nets (LLINs) and biological control may produce substantial transmission reduction in malaria transmitted by *An. culicifacies*^{90,91}. Most LLINs, however, employ pyrethroid insecticide against which already incipient resistance has been reported. There is an urgent need to develop innovative strategies against resistant *An. culicifacies* populations through research and development in key areas identified from field operations⁹²⁻⁹⁵.

Priority areas of research

An. culicifacies is the best studied mosquito species for its sibling species identification, range of distribution and relationship with malaria transmission in India. Yet, study of chromosomes (ovarian polytene chromosomes

for A, B, C, D, and mitotic chromosomes for B and E) is still the only technique that can distinguish sibling species A, B, C, D and E. No other probe distinguished each species unequivocally. PCR-based diagnostic assays distinguished only two distinct groups namely group I (species A/D) and group II (species B/C/E). Efforts to distinguish species of each group A/D and that of B/C/E based on PCR assays are still far from perfect and could not be used reliably⁵¹.

The present distribution of sibling species requires further research for understating population genetics structure in high risk areas of malaria. Recent findings revealed invasion by *An. culicifacies* sibling species A, D and E in addition to prevalent B and C in Odisha⁵⁴, and species 'E' in Madhya Pradesh⁵⁵. Similarly, in Assam (northeast India), species A is also observed to occur besides species B (Nanda N, personal communication). Additional investigations on distribution and bionomical characteristics of species E including insecticide susceptibility status, biting time to delimit its role in malaria transmission would be desirable in problem areas.

The crossing experiments between species A and B revealed post-mating barriers marked by unidirectional hybrid male sterility/ reduced fertility/ atrophied reproductive organs. Similar observations are deemed necessary to establish the extent of post-zygotic isolation by crossing experiments between other member sibling species. The inter-cross fertility data and the existence of possible morphological differences between sibling species A, B, C, D and E should be accorded priority to designate binomial nomenclature similar to other anopheline species vector taxa⁹⁶⁻⁹⁸.

An. culicifacies is highly adaptive, holds enormous behavioural plasticity and a fast invading species in areas hitherto found with low vector density, e.g. in deforested pockets in eastern and northeast India and breeding in altered ecological habitats^{23,99-101}, and believed to be undergoing significant genetic differentiation with obvious implications in its control¹⁰². There is a need to understand species-specific bionomics and insecticide susceptibility status before undertaking IRS for the control of *An. culicifacies* transmitted malaria.

The primary emphasis in the control of *An. culicifacies* should be on the innovative integrated methods that are cost-effective, safe and sustainable. It should, therefore, be realized that insecticides should be used sparingly and IRS when it becomes absolutely essential, e.g. epidemic situations. The routine vector control should rely on the bioenvironmental/ integrated

vector management methods. Field research should be undertaken on malaria control in various ecotypes such as the rural malaria, forest malaria, irrigation malaria, peri-urban malaria, industrial malaria, border malaria, coastal malaria by implementation of integrated vector management (IVM) methods involving the communities. The minimum area for the demonstration of new technologies should be a Primary Health Centre/ Community Health Centre and extended to cover a district. This would require mapping of malaria risk areas for *An. culicifacies* sibling species distribution, potential mosquito breeding habitats, site of contracting infection and control of outdoor malaria transmission. Insecticide incorporated plastic sheeting and insecticide treated hammock should be field tested in areas not responding to treated bed nets/LLINs^{103,104}.

Along with bionomics of *An. culicifacies*, socio-economic research should continue on human settlements, housing structure, vocations, sleeping habits, migration pattern, and community participation in vector control. Intensive information, education and communication (IEC) for various target groups, health system strengthening and capacity building including technology transfer to the control programme should constitute a continuing activity of the NVBDCP.

Conclusions

In the past decades, a wealth of data on the biology and control of *An. culicifacies* has been generated. *An. culicifacies* populations build up during monsoon and post monsoon season, and bring periodical focal outbreaks and epidemics throughout its range of distribution. Recent studies utilizing the molecular techniques for identification of sibling species complex have helped in understanding bionomics and role of each sibling species in malaria transmission. *An. culicifacies* is an invasive species and its further spread is supported by the changing environmental determinants. Therefore, entomological surveillance should be integral to malaria surveillance. *An. culicifacies* rapidly develops multiple insecticide resistance, and *inter alia* it is playing an increasing role in the transmission of *P. vivax* and *P. falciparum*, and spread of multi-drug resistant malaria. Research leading to newer evidence-based interventions, community based integrated vector management strategies should be strengthened in tackling the emerging challenges in the control of *An. culicifacies* transmitted malaria. The priority research areas in vector control may include developing malaria-risk maps, evidence based focused and selective vector control, tackling problems in vector control that

arise in the field, protection from contracting malaria in the settlements in forests including inaccessible areas, personal protection measures for individuals and communities, reduction in malaria receptivity legislative measures and biological control, vector surveillance, monitoring insecticide resistance and developing countermeasures, protecting target areas from vector invasion, cross-border synchronized vector control, and establishing priority in vector control to eliminate drug-resistant malaria foci.

References

- Singh V, Mishra N, Awasthi G, Dash AP, Das A. Why is it important to study malaria epidemiology in India? *Trends Parasitol* 2009; 25 : 452-7.
- Das A., Anvikar AR, Cator LJ, Dhiman RC, Eapen A, Mishra N, *et al.* Malaria in India: the centre for the study of complex malaria in India. *Acta Trop* 2012; 121 : 267-73.
- Sharma VP. Determinants of malaria in South Asia. In: Casman EA, Dowlatabadi H, editors. *The contextual determinants of malaria*. Pittsburgh PA: Carnegie Mellon University, USA; 2002. p.110-32.
- Malaria: magnitude of the problem. National Vector Borne Disease Control Programme. Ministry of Health & Family Welfare, Government of India. Available from: <http://www.nvbdc.gov.in>, accessed on April 6, 2015.
- Dhingra N, Jha P, Sharma VP, Cohen AA, Jotkar RM, Rodriguez PS, *et al.* Adult and child malaria mortality in India: a nationally representative mortality study. *Lancet* 2010; 376 : 1768-74.
- John TJ, Dandona L, Sharma VP, Kakkar M. Continuing challenge of infectious diseases in India. *Lancet* 2011; 377 : 252-69.
- Sharma VP. Hidden burden of malaria in Indian women. *Malar J* 2009; 8 : 281.
- Dev V, Sharma VP. The dominant mosquito vectors of human malaria in India. In: Manguin S, editor. *Anopheles mosquitoes - new insights into malaria vectors*. Croatia: InTech Publications; 2013. p. 239-71.
- Singh N, Mishra AK, Chand SK, Sharma VP. Population dynamics of *Anopheles culicifacies* and malaria in the tribal area of central India. *J Am Mosq Control Assoc* 1999; 15 : 283-90.
- Mathur KK, Harpalani G, Kalra NL, Murthy GG, Narasimham MV. Epidemic of malaria in Barmer district (Thar desert) of Rajasthan during 1990. *Indian J Malariol* 1992; 29 : 1-10.
- Bhuyan M, Das NG, Chakraborty BC, Talukdar PK, Sarkar PK, Das SC, *et al.* Role of *Anopheles culicifacies* during an outbreak of malaria in Garubandha P.H.C., Assam. *J Commun Dis* 1997; 29 : 243-6.
- Das NG, Gopalakrishnan R, Talukdar PK, Baruah I. Diversity and seasonal densities of vector anophelines in the relation to forest fringe malaria in district Sonitpur, Assam (India). *J Parasit Dis* 2011; 35 : 123-8.
- Barik TK, Sahu B, Swain V. A review on *Anopheles culicifacies*: from bionomics to control with special reference to Indian subcontinent. *Acta Trop* 2009; 109 : 87-97.
- Sharma VP. Re-emergence of malaria in India. *Indian J Med Res* 1996; 103 : 26-45.
- Akhtar R, Dutt AK, Wadhwa V, editors. Malaria in South Asia: eradication and resurgence during the second half of the twentieth century. *Advances in Asian human-environment research*. New York: Springer; 2010. p. 241.
- Nagpal BN, Sharma VP. *Indian anophelines*. New Delhi: Oxford & IBH Publishing; 1995. p. 416.
- World Health Organization (WHO). *Anopheles species complexes in South and South East-Asia*. SEARO Technical Publication No. 57. Geneva: WHO; 2007. p. 102.
- Rao TR. *The anophelines of India*. New Delhi: Malaria Research Centre, Indian Council of Medical Research; 1984. p. 596.
- Russell PF, Jacob VP. On the epidemiology of malaria in the Nilgiris district, Madras Presidency. *J Malar Inst India* 1942; 4 : 349-92.
- Jacob VP. Some aspects of malaria in Jammu and Kashmir state. *Indian J Malariol* 1950; 4 : 251-60.
- Rao TR, Dhanda V, Bhat HR, Kulkarni SM. A survey of haematophagous arthropods in Western Himalayas, Sikkim and hill districts of West Bengal. A general account. *Indian J Med Res* 1973; 61 : 1421-61.
- Covell G, Bailey JD. Malaria in Sind. Pts. 3. The factors influencing the normal autumnal incidence of malaria in Lakana Taluka, Larkana district. *Rec Mal Surv India* 1930; 1 : 549-65.
- Ramasamy R, Surendran SN. Global environment changes and salinity adaptation in mosquito vectors. Available from: <http://www.amazon.com/Environment-Changes-Salinity-Adaptation-Mosquito/dp/3848422905>, accessed on January 20, 2014.
- Ansari MA, Mani TR, Sharma VP. A preliminary note on the colonization of *Anopheles culicifacies* Giles. *J Commun Dis* 1977; 9 : 206-7.
- Green CA, Miles SJ. Chromosomal evidence for sibling species of the malaria vector *Anopheles (Cellia) culicifacies* Giles. *J Trop Med Hyg* 1980; 83 : 75-8.
- Subbarao SK, Vasantha K, Adak T, Sharma VP. *Anopheles culicifacies* complex: evidence for a new sibling species C. *Ann Entomol Soc Am* 1983; 76 : 985-8.
- Subbarao SK, Vasantha K, Sharma VP. Cytotaxonomy of certain malaria vectors of India. In: Service MW, editor. *Biosystematics of haematophagous insects*. Oxford: Clarendon Press; 1988. p. 25-37.
- Suguna SG, Tewari SC, Mani TR, Hiriyan J, Reuben R. A cytogenetic description of a new species of the *Anopheles culicifacies* complex. *Genetica* 1988-1989; 78 : 225-30.
- Vasantha K, Subbarao SK, Sharma VP. *Anopheles culicifacies* complex: population cytogenetic evidence for species D (Diptera: Culicidae). *Ann Entomol Soc Am* 1991; 84 : 531-6.
- Kar I, Subbarao SK, Eapen A, Ravindran J, Satyanarayana TS, Raghavendra K, *et al.* Evidence for a new malaria vector species, species E, within the *Anopheles culicifacies* complex (Diptera: Culicidae). *J Med Entomol* 1999; 36 : 595-600.
- Akoh JI, Beidas MF, White GB. Cytotaxonomic evidence for the malaria vector species A of the *Anopheles culicifacies* complex being endemic in Arabia. *Trans R Soc Trop Med Hyg* 1984; 78 : 698-700.

32. Zaim M, Subbarao SK, Manouchehri AV, Cochrane AH. Role of *Anopheles culicifacies* s. l. and *An. pulcherrimus* in malaria transmission in Ghassreghand (Baluchistan), Iran. *J Am Mosq Control Assoc* 1993; 9 : 23-6.
33. Mahmood F, Sakai RK, Akhtar K. Vector incrimination studies and observations on species A and B of the taxon *Anopheles culicifacies* in Pakistan. *Trans R Soc Trop Med Hyg* 1984; 78 : 607-16.
34. Abhayawardana TA, Dilrukshi RK, Wijesuriya SR. Cytotaxonomical examination of sibling species in the taxon *Anopheles culicifacies* Giles in Sri Lanka. *Indian J Malariol* 1996; 33: 74-80.
35. de Silva BG, Gunasekera MB, Abeyewickreme W, Karunanayake EH. Field analysis of *Anopheles culicifacies* Giles complex for the presence of sibling species A. *Proc Sri Lankan Assoc Adv Sci* 1993; 49 : 24.
36. de Silva BG. *Anopheles culicifacies* complex: geographical distribution of sibling species and existing methods for their identification. *Vidyodaya J Sci* 2009; 14 : 1-28.
37. Baimai V, Kijchalao U, Rattanarithikul R. Metaphase karyotypes of *Anopheles* of Thailand and Southeast Asia: V. The *Myzomyia* Series, subgenus *Cellia* (Diptera: Culicidae). *J Am Mosq Control Assoc* 1996; 12 : 97-105.
38. Miles SJ. Unidirectional hybrid male sterility from crosses between species A and species B of the taxon *Anopheles (Cellia) culicifacies* Giles. *J Trop Med Hyg* 1981; 84 : 13-6.
39. Subbarao SK, Vasantha K, Sharma VP. Studies on the crosses between the sibling species of the *Anopheles culicifacies* complex. *J Hered* 1988; 79 : 300-3.
40. Vasantha K, Subbarao SK, Adak T, Sharma VP. Karyotypic variations in *Anopheles culicifacies* complex. *Indian J Malariol* 1982; 19 : 27-32.
41. Vasantha K, Subbarao SK, Adak T, Sharma VP. *Anopheles culicifacies*: mitotic karyotype of species C. *Indian J Malariol* 1983; 20 : 161-2.
42. Suguna SG, Tewari SC, Mani TR, Hiriyani J, Reuben R. *Anopheles culicifacies* species complex in Thenpennaiyar riverine tract, Tamil Nadu. *Indian J Med Res* 1983; 77 : 455-9.
43. Subbarao SK, Nanda N, Chandras RK, Sharma VP. *Anopheles culicifacies* complex: cytogenetic characterization of Rameshwaram islands populations. *J Am Mosq Control Assoc* 1993; 9 : 27-31.
44. Adak T, Subbarao SK, Sharma VP, Rao SR. Lactate dehydrogenase allozyme differentiation of species in the *Anopheles culicifacies* complex. *Med Vet Entomol* 1994; 8 : 137-40.
45. Milligan PJ, Phillips A, Molyneux DH, Subbarao SK, White GB. Differentiation of *Anopheles culicifacies* Giles (Diptera: Culicidae) sibling species by analysis of cuticular components. *Bull Entomol Res* 1986; 76 : 529-37.
46. Gunasekera MB, de Silva BG, Abeyewickreme W, Subbarao SK, Nandadasa HG, Karunanayake EH. Development of DNA probes for the identification of sibling species A of the *Anopheles culicifacies* (Diptera: Culicidae) complex. *Bull Entomol Res* 1995; 85 : 345-53.
47. Singh OP, Goswami G, Nanda N, Raghavendra K, Chandra D, Subbarao SK. An allele-specific polymerase chain reaction assay for the differentiation of members of the *Anopheles culicifacies* complex. *J Biosci* 2004; 29 : 275-80.
48. Goswami G, Raghavendra K, Nanda N, Gakhar SK, Subbarao SK. PCR-RFLP of mitochondrial cytochrome oxidase subunit II and ITS2 of ribosomal DNA: markers for the identification of members of the *Anopheles culicifacies* complex (Diptera: Culicidae). *Acta Trop* 2005; 95 : 92-9.
49. Manonmani AM, Sadanandane C, Sahu SS, Mathivanan A, Jambulingam P. rDNA-ITS2-PCR assay for grouping the cryptic species of *Anopheles culicifacies* complex (Diptera: Culicidae). *Acta Trop* 2007; 104 : 72-7.
50. Goswami G, Singh OP, Nanda N, Raghavendra K, Gakhar SK, Subbarao SK. Identification of all members of the *Anopheles culicifacies* complex using allele-specific polymerase chain reaction assays. *Am J Trop Med Hyg* 2006; 75 : 454-60.
51. Surendran SN, Hawkes NJ, Steven A, Hemingway J, Ramasamy R. Molecular studies of *Anopheles culicifacies* (Diptera: Culicidae) in Sri Lanka: sibling species B and E show sequence identity at multiple loci. *Eur J Entomol* 2006; 103 : 233-7.
52. Raghavendra K, Cornel AJ, Reddy BP, Colins FH, Nanda N, Chandra D, et al. Multiplex PCR assay and phylogenetic analysis of sequences derived from D2 domain of 28S rDNA distinguished members of the *Anopheles culicifacies* complex into two groups, A/D and B/C/E. *Infect Genet Evol* 2009; 9 : 271-7.
53. Tripathy A, Samanta L, Das S, Paride SK, Marai N, Hazra RK, et al. Distribution of sibling species of *Anopheles culicifacies* s.l. and *Anopheles fluviatilis* s.l. and vectorial capacity in eight different malaria endemic districts of Orissa, India. *Mem Inst Oswaldo Cruz* 2010; 105 : 981-7.
54. Das M, Das B, Patra AP, Tripathy HK, Mohapatra N, Kar SK, et al. *Anopheles culicifacies* sibling species in Odisha, Eastern India: First appearance of *Anopheles culicifacies* E and its vectorial role in malaria transmission. *Trop Med Int Health* 2013; 18 : 810-21.
55. Sharma AK, Tyagi V, Singh S, Veer V, Agrawal OP, Sukumaran D. Distribution of *Anopheles culicifacies* and detection of its sibling species E from Madhya Pradesh: Central India. *J Arthropod-Borne Dis* 2014; 8 : 182-96.
56. Subbarao SK, Vasantha K, Adak T, Sharma VP. Seasonal prevalence of sibling species A and B of the taxon *Anopheles culicifacies* in villages around Delhi. *Indian J Malariol* 1987; 24 : 9-15.
57. National Institute of Malaria Research (NIMR). *Anopheles culicifacies* and *An. fluviatilis* complexes and their control. Technical Report series No. NIMR/TRS/2009-January/2002. New Delhi, India: NIMR; 2002.
58. Subbarao SK, Adak T, Vasantha H, Joshi H, Raghavendra K, Cochrane AH, et al. Susceptibility of *Anopheles culicifacies* species A and B to *Plasmodium vivax* and *Plasmodium falciparum* as determined by immunoradiometric assay. *Trans R Soc Trop Med Hyg* 1988; 82 : 394-7.
59. Subbarao SK, Sharma VP. Anopheline species complexes & malaria control. *Indian J Med Res* 1997; 106 : 164-73.
60. Subbarao SK, Nanda N, Raghavendra K. Malariogenic stratification of India using *Anopheles culicifacies* sibling species prevalence. *ICMR Bull* 1999; 29 : 75-80.
61. Sharma A, Parasher H, Singh OP, Adak T. Species B of *Anopheles culicifacies* (Diptera: Culicidae) is reproductively less fit than species A and C of the complex. *Acta Trop* 2009; 112 : 316-9.

62. Adak T, Kaur S, Singh OP. Comparative susceptibility of different members of the *Anopheles culicifacies* complex to *Plasmodium vivax*. *Trans R Soc Trop Med Hyg* 1999; 93 : 573-7.
63. Kaur S, Singh OP, Adak T. Susceptibility of species A, B, and C of *Anopheles culicifacies* complex to *Plasmodium yoelii yoelii* and *Plasmodium vinckei petteri* infections. *J Parasitol* 2000; 86 : 1345-8.
64. Subbarao SK, Vasantha K, Sharma VP. Responses of *Anopheles culicifacies* sibling species A and B to DDT and HCH in India: implications in malaria control. *Med Vet Entomol* 1988; 2 : 219-23.
65. Raghavendra K, Subbarao SK, Vasantha K, Pillai MK, Sharma VP. Differential selection of malathion resistance in *Anopheles culicifacies* A and B (Diptera: Culicidae) in Haryana state, India. *J Med Entomol* 1992; 29 : 183-7.
66. Tiwari SN, Prakash A, Subbarao SK, Roy A, Joshi H, Sharma VP. Correlation of malaria endemicity with *An. culicifacies* sibling species composition and malaria antibody profile in distort Allahabad (U.P.). *Indian J Malariol* 1994; 31 : 48-56.
67. Sharma VP, Sharma RC, Gautam AS. Bio-environmental control of malaria in Nadiad, Kheda district, Gujarat. *Indian J Malariol* 1986; 23 : 95-117.
68. Sharma VP. Community-based malaria control in India. *Parasitol Today* 1987; 3 : 222-6.
69. Sharma VP, Ghosh A, editors. Larvivorous fishes of inland ecosystems. *Proceedings of the MRC-CICFRI Workshop*. Delhi: Malaria Research Centre (ICMR); 1994. p. 224.
70. Ghosh SK, Tewari SN, Sathyanarayan TS, Sampath TR, Sharma VP, Nanda N, et al. Larvivorous fish in wells target the malaria vector sibling species of the *Anopheles culicifacies* complex in vaillages in Karnataka, India. *Trans R Soc Trop Med Hyg* 2004; 99 : 101-5.
71. Singh N, Mehra RK, Srivastava N. Malaria during pregnancy and infancy, in an area of intense malaria transmission in central India. *Ann Trop Med Parasitol* 2001; 95 : 19-29.
72. Nanda N, Yadav RS, Subbarao SK, Joshi H, Sharma VP. Studies on *Anopheles fluviatilis* and *Anopheles culicifacies* sibling species in relation to malaria in forested hilly and deforested riverine ecosystems in northern Orissa, India. *J Am Mosq Control Assoc* 2000; 16 : 199-205.
73. Dutt AK, Dutta HM, Parera C. Resurgence of malaria in Sri Lanka in the 1970s. In: Akhtar R, Dutt AK, Wadhwa V, editors. *Malaria in South Asia: eradication and resurgence during the second half of the twentieth century*. New York: Springer; 2010. p. 29-41.
74. Zaidi IH, Kazmi JH. The resurgence of malaria in Pakistan: a geographical evaluation. In: Akhtar R, Dutt AK, Wadhwa V, editors. *Malaria in South Asia: eradication and resurgence during the second half of the twentieth century*. New York: Springer; 2010. p. 123-39.
75. Singh N, Singh OP, Sharma VP. Dynamics of malaria transmission in forested and deforested regions of Mandla district, Central India (Madhya Pradesh). *J Am Mosq Control Assoc* 1996; 12 : 225-34.
76. Singh N, Chand SK, Bharti PK, Singh MP, Chand G, Mishra AK, et al. Dynamics of forest malaria transmission in Balaghat district, Madhya Pradesh, India. *PLoS One* 2013; 8 : e73730.
77. Raghavendra K, Vasantha K, Subbarao SK, Pillai MK, Sharma VP. Resistance in *Anopheles culicifacies* sibling species B and C to malathion in Andhra Pradesh and Gujarat States, India. *J Am Mosq Control Assoc* 1991; 7 : 255-9.
78. Singh OP, Raghavendra K, Nanda N, Mittal PK, Subbarao SK. Pyrethroid resistance in *Anopheles culicifacies* in Surat district, Gujarat, India. *Curr Sci* 2002; 82 : 547-50.
79. Bhatt RM, Sharma SN, Barik TK, Raghavendra K. Status of insecticide resistance in malaria vector, *Anopheles culicifacies* in Chhattisgarh state, India. *J Vector Borne Dis* 2012; 49 : 36-8.
80. Mishra AK, Chand SK, Barik TK, Dua VK, Raghavendra K. Insecticide resistance status in *Anopheles culicifacies* in Madhya Pradesh, Central India. *J Vector Borne Dis* 2012; 49 : 39-41.
81. Singh OP, Bali P, Hemingway J, Subbarao SK, Dash AP, Adak T. PCR-based methods for the detection of L1014 *kdr* mutation in *Anopheles culicifacies sensu lato*. *Malar J* 2009; 8 : 154.
82. Hoti SL, Vasuki V, Jambulingam P, Sahu SS. *kdr* allele-based PCR assay for detection of resistance to DDT in *Anopheles culicifacies sensu lato* Giles population from Malkangiri District, Orissa, India. *Curr Sci* 2006; 91 : 658-61.
83. Sharma VP, Prasittisuk C, Kondrashin AV. Magnitude of forest related malaria in the WHO South East Asia region. In: Sharma VP, Kondrashin AV, editors. *Forest malaria in South East Asia. Proceedings of an informal consultative meeting of WHO/MRC*. 1991 Feb 18-22; New Delhi: Malaria Research Centre (ICMR); 1991. p. 29-53.
84. Kalra NL. Forest malaria vectors in India: ecological characteristics and epidemiological implications. In: Sharma VP, Kondrashin AV, editors. *Forest malaria in South East Asia. Proceedings of an informal consultative meeting of WHO/MRC* 1991. Feb.18-22; New Delhi: Malaria Research Centre; 1991. p. 93-114.
85. World Health Organization. *Global plan for insecticide resistance management in malaria vectors (GPIRM)*. Available from: http://www.who.int/malaria/vector_control/ivm/gpirm, accessed on July 2, 2012.
86. Sharma SK, Upadhyay AK, Haque MA, Tyagi PK, Kindo BK. Impact of changing over of insecticide from synthetic pyrethroids to DDT for indoor residual spray in a malaria endemic area of Orissa, India. *Indian J Med Res* 2012; 135 : 382-8.
87. Raghavendra K, Barik TK, Reddy BP, Sharma P, Dash AP. Malaria vector control: from past to future. *Parasitol Res* 2011; 108 : 757-79.
88. Sharma RC, Gautam AS, Bhatt RM, Gupta DK, Sharma VP. The Kheda malaria project: the case for environmental control. *Health Policy Plann* 1991; 6 : 262-70.
89. Dev V, Sharma VP, Hojai D. Malaria transmission and disease burden in Assam: challenges and opportunities. *J Parasit Dis* 2009; 33 : 13-22.
90. Sreehari U, Mittal PK, Razdan RK, Ansari MA, Rizvi MM, Dash AP. Efficacy of PermaNet 2.0 against *Anopheles culicifacies* and *Anopheles stephensi* malaria vectors in India. *J Am Mosq Control Assoc* 2007; 23 : 220-3.
91. Sharma SK, Upadhyay AK, Haque MA, Tyagi PK, Raghavendra K, Dash AP. Wash-resistance and field evaluation of alphacypermethrin treated long-lasting insecticidal net (Interceptor) against malaria vectors, *Anopheles culicifacies*

- and *Anopheles fluviatilis* in a tribal area of Orissa, India. *Acta Trop* 2010; 116 : 24-30.
92. Sharma VP. Battling malaria iceberg incorporating strategic reforms in achieving Millennium Development Goals & malaria elimination in India. *Indian J Med Res* 2012; 136 : 907-25.
 93. Bhatia R, Rastogi RM, Ortega L. Malaria successes and challenges in Asia. *J Vector Borne Dis* 2013; 50 : 239-47.
 94. Pannetier C, Bouraima A, Chandre F, Piumeu M, Etang J, Rossignol M, et al. Efficacy of Olyset Plus, a new long-lasting insecticidal net incorporating permethrin and piperonyl butoxide against multi-resistant malaria vectors. *PLoS One* 2013; 8 : e75134.
 95. Corbel V, Chabi J, Dabire RK, Etang J, Nwane P, Pigeon O, et al. Field efficacy of a new mosaic long lasting mosquito net (PermaNet 30) against pyrethroid-resistant malaria vectors: a multi centre study in Western and Central Africa. *Malar J* 2010; 9 : 113.
 96. Sallum MA, Peyton EL, Wilkerson RC. Six new species of the *Anopheles leucosphyrus* group, reinterpretation of *An. elegans* and vector implications. *Med Vet Entomol* 2005; 19 : 158-99.
 97. Manguin S, Garros C, Dusfour I, Harbach RE, Coosemans M. Bionomics, taxonomy, and distribution of the major malaria vector taxa of *Anopheles* subgenus *Cellia* in Southeast Asia: an updated review. *Infect Genet Evol* 2008; 8 : 489-503.
 98. Sinka ME, Bangs MJ, Manguin S, Chareonviriyaphap T, Patil AP, Temperley WH, et al. The dominant *Anopheles* vectors of human malaria in the Asia-Pacific region: occurrence data, distribution maps and bionomic précis. *Parasit Vectors* 2011; 4 : 89.
 99. Gunathilaka N, Fernando T, Hapugoda M, Wickremasinghe R, Wijeyerathne P, Abeyewickreme W. *Anopheles culicifacies* breeding in polluted water bodies in Trincomalee district of Sri Lanka. *Malar J* 2013; 12 : 285.
 100. Sharma SK, Tyagi PK, Padhan K, Upadhyay AK, Haque MA, Nanda N, et al. Epidemiology of malaria transmission in forest and plain ecotype villages in Sundargarh district, Orissa, India. *Trans R Soc Trop Med Hyg* 2006; 100 : 917-25.
 101. Tyagi V, Sharma AK, Yadav R, Agrawal OP, Devanathan S, Veer V. Characteristics of the larval breeding sites of *Anopheles culicifacies* sibling species in Madhya Pradesh, India. *Int J Malar Res Rev* 2013; 1 : 47-53.
 102. Sunil S, Singh OP, Nanda N, Raghavendra K, Reddy BP, Subbarao SK. Analysis of population genetic structure of Indian *Anopheles culicifacies* species A using microsatellite markers. *Parasit Vectors* 2013; 6 : 166.
 103. Chandre F, Dabire RK, Hougard JM, Djogbenou LS, Irish SR, Rowland M, et al. Field efficacy of pyrethroid treated plastic sheeting (durable lining) in combination with long lasting insecticidal nets against malaria vectors. *Parasit Vectors* 2010; 3 : 65.
 104. Grietens KP, Xuan XN, Ribera J, Duc TN, Bortel Wv, Ba NT, et al. Social determinants of long lasting insecticidal hammock use among the Ra-glai ethnic minority in Vietnam: implications for forest malaria control. *PLoS One* 2012; 7 : e29991.

This write-up contributed by Dr. V.P. Sharma, Centre for Rural Development and Technology, Indian Institute of Sciences, New Delhi (Formerly Additional Director General, ICMR, New Delhi) and Dr. Vas Dev, Scientist F, National Institute of Malaria Research Field Station, Guwahati, has been adapted from Indian Journal of Medical Research, Vol. 141, May 2015. pp.525-536.